

Test Hays for Nutrient Composition Before Feeding

Shane Gadberry
Associate Professor -
Animal Science

Mark Keaton
County Extension Agent -
Staff Chair

Variation in Hay Quality

Developing a least-cost, supplemental feeding program begins by having hays analyzed for nutrient content. A forage analysis is necessary because hays can be highly variable in nutrient content. Bermudagrass samples submitted to the University of Arkansas Agricultural Diagnostic Lab illustrate this point (Figure 1). Bermudagrass hays ranged from 6 to 22 percent protein. Utilization of the Arkansas average protein content of 13 percent for bermudagrass could result in under-feeding protein in some situations.





Figure 1. Distribution of crude protein content of bermudagrass hay samples submitted to the UA Agricultural Diagnostic Service Laboratory.

An evaluation of hay samples collected over a five-year period from farms participating in the Arkansas Beef Improvement Program (ABIP) indicates that hay quality can be highly variable from year to year. A statistical measure often used for quality control is called coefficient of variation (CV). Large CVs indicate a lot of variation or poor consistency.

Coefficients of variation smaller than 10 are desirable. Arkansas Beef Improvement Program data indicates protein may be more variable than fiber. Ninety percent of the farms had a CV greater than 10 percent for protein, and 40 percent of the farms had a CV greater than 10 percent for acid detergent fiber, which affects the digestibility of forages. The high variability in hays across the state, even among hays produced on a single farm, makes annual hay testing important.

Tips for Collecting Forage Samples

The analyzed nutrient composition of hay is only as accurate as the sample sent. Begin by accurately identifying the dominant forage species of the field harvested to get an accurate estimate of total digestible nutrients (TDN).



Test each lot of hay. A lot represents hay harvested from each cutting within a single field under similar harvest conditions. Harvest date, plant species,

*Arkansas Is
Our Campus*

Visit our web site at:
<http://www.uaex.edu>

fertility and weather can all result in variations within and between fields and cuttings. If testing each lot of hay is impractical, testing across hay lots is the next best approach. For mixed lot sampling, the combined sample should contain each lot's proportion relative to the whole. If lot 1 contains 15 percent of the total number of bales, then it should also represent 15 percent of mixed lot sample. Otherwise, forage quality will be interpreted inaccurately. For example, sampling the same number of bales from lots 1, 2 and 3 (Table 1) would result in an overestimate of TDN and result in under-feeding beef cattle during early and mid-lactation.

Table 1. Effect of Sampling Across Hay Lots on Total Digestible Nutrients

Lot	Number of Bales	Percent of Total Bales	Sample Size Bales/Lot (30%)	Total Digestible Nutrients
1	30	15%	9	64%
2	30	15%	9	58%
3	140	70%	42	52%
Non-weighted Average ¹				58.0%
Weighted Average ²				54.7%

¹Non-weighted average TDN reflects an inaccurate estimate of TDN that could occur if the same number of bales were sampled from each lot.

²Weighted average TDN reflects a more accurate estimate of overall hay TDN because the hay sample is proportional to the size of each lot [(64 x 0.15) + (58 x 0.15) + (52 x 0.70)].

Core samples are preferred over grab samples. Even with hay that is not weathered, multiple core samples will contain a better distribution of plant material, which will result in a more accurate assessment of nutrient composition. Hay sampling probes (Figure 2) are available for use through county Extension offices in Arkansas. Samples should be

taken from the end of square bales and from the side of round bales and stacks.



Figure 2. Core sampling devices: (1) Penn State Forage sampler, (2) Colorado Hay Probe and (3) Star Quality Multi-sampler.

Twenty to thirty percent of the bales must be sampled to accurately estimate the nutrient composition of the hay. A demonstration project at the University of Arkansas showed differences of 5 percent TDN when as few as 5 percent of bales were sampled within a single hay lot (Figure 3). A minimum of six individually core-sampled round bales are necessary to have sufficient sample size for an analysis. Sample size should represent the larger of the two – either 6 bales or 20 percent of the number of bales in a lot.




Figure 3. Effect of sample size on bermudagrass TDN (Milliken et al., Ark. Animal Sci., 2003).

Hay that is stored outside should be sampled just prior to the feeding period. Angle the core sampling tool in an upward direction when sampling large packages stored outside. This will help avoid creating a hole for water to penetrate the inside of the bale. Avoid sampling excessively weathered material. The sample submitted should reflect what the cattle will be eating, not wasting. Hay that is stored inside can be sampled at storage or prior to feeding, with the exception of hay that was harvested wet. Hay quality diminishes rapidly during storage if harvested at less than 80 percent dry matter.

Using Results

Upon receiving the results from a forage analysis, the next step is to interpret the results. Figure 4 is an example routine hay analysis report. The results are separated into two columns – **AS FED BASIS** and **DRY MATTER BASIS**. When comparing a forage analysis to animal requirements, the values reported under dry matter basis should be used. The following information is reported when a forage sample is submitted for routine analysis through the University of Arkansas Cooperative Extension Service.

- **Moisture** – water content of a feed
- **Dry Matter** – 100 minus the water content. Used to convert AS FED to DRY MATTER. Example: $8.9 - 91.3 \times 100 = 9.8$.
- **Crude Protein** – a measure of plant nitrogen multiplied times 6.25.
- **Acid Detergent Fiber** – a measure of plant cellulose and lignin. Acid detergent fiber is commonly used to estimate digestibility (TDN).
- **Neutral Detergent Fiber** – a measure of plant hemicellulose, cellulose and lignin. Neutral detergent fiber may also be used to estimate digestibility and is highly correlated with forage intake.
- **Total Digestible Nutrients (TDN)** – an estimate of the supply of energy. The number for total digestible nutrients is derived from equations developed from feeding trials. Equations commonly utilize acid detergent fiber to estimate TDN but may also include crude protein and/or neutral detergent fiber.
- **Net Energy for Lactation** – an estimate of the supply of energy commonly used for balancing dairy cow rations.

Agricultural Diagnostic Service Laboratory
University of Arkansas
Crop, Soil, and Environmental Sciences

FEED ANALYSIS REPORT

(STATE COPY)



ANALYSIS FOR :

Name-----	Date Mailed-- 7 July 2006
Address--	
City-----	State-- AR Zip-- 71832
County--	

Lab #-----F 60308
Date Received-- 07/03/06
Sample ID----- HOYT
Species Name-- MIXED GRASS
Species #----- 13
Variety-----
Type of sample- HAY
Analysis Date-- 07/07/06
Payment Received-# 9051

CHEMICAL COMPOSITION :

	AS FED BASIS	DRY MATTER BASIS
Moisture-----	8.7 %	
Dry Matter-----	91.3 %	
Crude Protein-----	8.9 %	9.8 %
Acid Detergent Fiber----	33.9 %	37.1 %
Neutral Detergent Fiber-	59.6 %	65.3 %

NUTRIENT AND ENERGY VALUE :

	AS FED BASIS	DRY MATTER BASIS
TOTAL DIGESTIBLE NUTRIENTS (TDN)--	48.6 %	53.2 %
NET ENERGY FOR LACTATION (NEL)----	0.49 Mcal/cwt	0.54 Mcal/cwt

ADDITIONAL:

Figure 4. Example routine analysis results summary prepared by the University of Arkansas Agricultural Diagnostic Service Laboratory.

Table 2. Nutrient Requirements of a Mature Beef Cow (adapted from *Beef Cattle Nutrition Series, Part 2: Establishing Nutritional Requirements, FSA3079*)

Group	CP % DM	TDN % DM
1100 lb cow, early lactation (20 lb peak milk)	11	60
1100 lb cow, mid-lactation	9	56
1100 lb cow @ calf weaning, mid-gestation	7	47
1100 lb cow @ late gestation	8	52

A forage test cannot be interpreted without having a set of livestock nutritional requirements to compare against the results. Nutrient requirement tables are available through the county Extension office for many classes of livestock. Table 2 contains the nutrient requirements for a mature beef cow during different stages of production. Compared to the example forage test, the hay quality is sufficient to meet the nutritional requirements of both a cow in mid- and late gestation. A cow in early lactation would require both supplemental protein and energy (TDN) because a lactating cow's requirement exceeds the amount in the hay, and a cow in mid-lactation would only require supplemental energy.

Table 3. Percent of Ozark Bootstrap Samples Meeting Requirements for Different Classes of Beef Cattle

Group	% Adequate
Weaning, mid-gestation	100
Late gestation	96
Mid-lactation	69
Early lactation	34

Baxter, Boone, Marion and Searcy counties in Arkansas reported approximately 34 percent of the hay samples collected (Table 3) from 23 farms participating in the Ozark Bootstrap (pilot multi-county)

demonstration project were sufficient to meet the nutritional requirements of a mature beef cow in early lactation. Nearly all of the forage samples (96 percent) met the protein and TDN requirement for a mature beef cow in late gestation.

Hay is usually harvested at a mature stage, which results in a shortage in TDN for cows in early lactation. Forage test results were used to improve the economics of winter feeding on ABIP and Bootstrap demonstration farms by:

- Matching hay lots with nutritional requirements of cows (feeding the best quality hay to lactating cows and lower-quality hay to gestating cows).
- Identifying which nutrient(s) were inadequate relative to beef cattle requirements and supplementing accordingly. Seventy percent of the hay samples from the Ozark Bootstrap demonstration **were** capable of meeting or exceeding the protein requirement of a mature beef cow in early lactation. However, 70 percent of the hay samples **were not** capable of meeting the energy requirement of a beef cow during early lactation. As a result, supplemental feeding was shifted from emphasizing protein to emphasizing energy.

Summary

The time and money spent forage testing has consistently been shown to be a valuable tool for avoiding costly feeding errors. However, a forage test is of little value if the producer is not willing to interpret the results and make supplemental feeding changes when necessary.

Visit the Animal Science Blog at www.arkansas-livestock.com.

Printed by University of Arkansas Cooperative Extension Service Printing Services.

DR. SHANE GADBERRY is associate professor - animal science, Department of Animal Science, Little Rock. **MARK KEATON** is county Extension agent - staff chair, Baxter County. They are employees of the University of Arkansas Division of Agriculture.

FSA3114-PD-12-12RV

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.