

**ATRAZINE AND WATER QUALITY**

Atrazine label restrictions regarding mixing, loading and application are discussed below. These restrictions are part of the overall ground and surface water contamination risk reduction measures. Atrazine users are strongly encouraged to follow these guidelines to comply with the label, and to share in the responsibility of preserving the future of this extremely valuable corn herbicide. **These restrictions, and the Restricted Use Pesticide designation, apply to all formulations of atrazine, and all package mix products which contain atrazine.**

**Mixing, Loading and Application** – Atrazine may not be mixed/loaded or used within 50 feet of all wells, including abandoned wells, drainage wells and sink holes. Atrazine may not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Atrazine may not be applied aerially or by ground within 66 feet of the points where field surface runoff enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If atrazine is applied to highly erodible land, the 66-foot buffer or setback from runoff entry points must be planted to corn, seeded with grass, or another suitable crop.

**Application rates – All soil applications prior to crop emergence –**

**\*Highly Erodible Soils** (as defined by NRCS) – If conservation tillage is practiced (at least 30 percent of residue coverage at planting), apply a maximum of 2 lb a.i./acre. If residue coverage is less than 30 percent, apply a maximum of 1.6 lb a.i./acre.

**\*Soils Not Highly Erodible** – Apply a maximum of 2 lb a.i./acre.

**Postemergence Applications** – If no atrazine was applied prior to corn emergence, apply a maximum of 2 lb a.i./acre. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lb a.i./acre/calendar year. Postemergence application to corn must be made before corn exceeds 12 inches in height.

<b>Crop, Situation, and Active Chemical Per Broadcast Acre</b>	<b>Weeds Controlled</b>	<b>Formulated Material Per Broadcast Acre</b>	<b>Time of Application</b>	<b>Method of Application and Precautions</b>
<b>FIELD CORN</b>				
<b>Removing Partial Corn Stands for Replant</b>				
clethodim @ 0.045 lb/A	Corn, including Roundup Ready and Liberty Link.	<b>Select Max</b> 0.97 lb/gal, 6 oz/A.	Up to 12-inch corn. Use crop oil concentrate + AMS.	Do not plant corn for 7 days after application.
glufosinate @ 0.6 lb/A	Corn, including Roundup Ready but not Liberty Link.	<b>Glufosinate (280 formulations)</b> 32 oz/A.	Good coverage is essential. Adding AMS may enhance control.	Note that corn with the Herculex insecticide trait is tolerant to glufosinate and will not be controlled. Corn may be replanted immediately.
paraquat + metribuzin @ 0.625 + 0.14 lb/A	Corn, including Roundup Ready and Liberty Link.	<b>Paraquat (2 or 3 lb/gal formulations) + metribuzin 75DF</b> 40 or 26 oz/A + 3 oz/A.	Good coverage is essential.	Corn may be replanted immediately.
paraquat + diuron @ 0.625 + 0.5 lb/A	Corn, including Roundup Ready and Liberty Link.	<b>Paraquat (2 or 3 lb/gal formulations) + Direx 4L</b> 40 or 26 oz/A + 1 pt/A.	Good coverage is essential.	Corn may be replanted immediately.
paraquat + atrazine @ 0.625 + 0.5 lb/A	Corn, including Roundup Ready and Liberty Link.	<b>Paraquat (2 or 3 lb/gal formulations) + Atrazine 4L</b> 40 or 26 oz/A + 1 pt/A.	Good coverage is essential.	Corn may be replanted immediately.
<b>Preemergence</b>				
Many herbicide “pre-mixes” exist for field corn – too many to list here. Check the active ingredients and rates for all pre-mixes. Most individual components of these mixes are listed below and in the weed control table (page 68).				
S-metolachlor @ 0.75 to 1.3 lb/A	Annual grasses and pigweed.	<b>Dual II Magnum 7.64 EC</b> 0.8 to 1.4 pt/A.	Preplant to preemerge.	
atrazine @ 2 lb/A	Most small-seeded annuals, annual morningglory, cocklebur, velvetleaf, smartweed and sicklepod.	<b>AAtrex, Atrazine</b> 2.5 lb/A 80W or 2 qt/A 4L or 2.2 lb/A Nine-0.	At planting.	Do not plant fall cover crops. Do not plant crops other than corn or grain sorghum in treated fields during the same season. Do not apply more than 2.5 lb/A active atrazine per season.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
atrazine @ 1.5 to 2.25 lb/A + 1 to 1.6 lb/A	Annual grasses, pigweed, annual morningglory, common cocklebur, velvetleaf, smartweed and sicklepod.	<b>Micro-Tech 4L + AAtrex, Atrazine</b> 1.5 to 2.25 qt/A Lasso + 1.25 to 2 lb/A 80W or 1 to 1.6 qt/A 4L or 1.1 to 1.75 lb/A Nine-0. or <b>Lariat/Bullet 4F</b> 2.5 to 3.75 qt/A.	Preemergence or preplant. <b>Note</b> – This treatment can be applied to emerged corn before it exceeds 5" tall. However, weed control will be reduced if weeds exceed 2" tall.	Add additional atrazine for improved control of cocklebur and morningglory. Rainfall in 5 to 7 days is necessary for best results. With preplants, shallow incorporate 2 to 3 inches within 7 days of planting.
S-metolachlor + atrazine @ 0.75 to 1.3 lb/A + 1 to 1.6 lb/A	Annual grasses, pigweed, annual morningglory, common cocklebur, velvetleaf, smartweed and sicklepod.	<b>Cinch or Dual II Magnum 7.64 EC + AAtrex, Atrazine</b> See label for specific formulations in question. 0.8 to 1.4 pt/A + 1.25 lb/A 80W or 2 pt/A 4L to 2.0 lb 80W or 3.2 pt/A 4L. or <b>Bicep II Magnum 5.5 L or Cinch ATZ</b> 1.3 to 2 qt/A.	Preemergence or preplant.	Add additional atrazine for improved control of cocklebur and morningglory. Rainfall in 5 to 7 days is necessary for best results. With preplants, shallow incorporate 2 to 3 inches within 7 days of planting. If concerned about achieving a stand, leave atrazine out as pre and follow with atrazine early post.
dimethenamid + atrazine @ 0.56 to 0.75 lb/A + 0.75 to 2 lb/A	Annual grasses, pigweed, annual morningglory, common cocklebur, velvetleaf and smartweed.	<b>Outlook 6E + AAtrex, Atrazine</b> 12 to 16 + 0.75 to 2 qt/A 4L	From 45 days preplant to pre-emergence up to 8" tall corn.	Same as above. Rates depend on percent organic matter. See label.
dimethenamid + saflufenacil @ 0.31 to 0.62 + 0.044 to 0.088 lb/A	Annual grasses, pigweed, velvetleaf, morningglory and horseweed.	<b>Verdict</b> 10 to 12 oz/A.	Burndown up to preemergence. <b>Do not</b> apply Verdict over the top of emerged corn.	Rainfall or overhead irrigation is required for activation. Verdict can be used as a burndown that leaves behind residual control. For best burndown activity, tank mix with glyphosate and use MSO 1 pt/A + AMS. On medium to fine soils, the rotation interval to soybeans is 30 days if you were to lose the corn crop. See label for restrictions.
acetochlor @ 1.09 to 1.97 lb/A	Annual grasses and pigweed.	<b>Surpass NXT</b> 1.25 to 2.25 pt/A.	Preplant or preemergence.	
acetochlor + atrazine @ 1.7 lb/A + 0.8 lb/A	Annual grasses, pigweed, morningglory, cocklebur, velvetleaf, smartweed and sicklepod.	<b>Degree 3.8 SL + Atrazine</b> 3.5 pt + 0.8 qt/A Atrazine 4L. or <b>Degree Xtra</b> 5 pt/A.	Preplant or preemergence.	Add additional atrazine for improved control of cocklebur and morningglory. Rainfall in 5 to 7 days is necessary for best results. With preplants, shallow incorporate 2 to 3 inches within 7 days of planting.
metolachlor + mesotrione + bicyclopyrone + atrazine @ 1.34 + 0.148 + 0.037 + 0.624 lb/A	Annual grasses, yellow nutsedge and broadleaf weeds.	<b>Acuron</b> 2.5 qt/A.	Preplant or preemergence.	May be applied up to 28 days before planting. Do not plant crops other than corn in treated area.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
<b>FIELD CORN</b>				
<b>Preemergence [cont.]</b>				
acetochlor + atrazine @ 0.85 to 2 lb/A + 0.88 to 1.6 lb/A	Annual grasses, pigweed, morningglory, common cocklebur, velvetleaf, smartweed and sicklepod.	<b>Keystone NXT</b> 1.4 to 2.6 qt/A.	Preplant or preemergence.	Add additional atrazine for improved control of morningglory.
acetochlor + clopyralid + flumetsulam @ 0.7 to 1.4 + 0.07 to 0.14 + 0.023 to 0.045 lb/A	Annual grasses and broadleaves, thistles.	<b>Surestart II</b> 1.5 to 2.5 pt/A.	Preplant or preemergence.	Optimal weed control will be obtained when applications are as close as possible to planting but before weeds emerge. Applications may be made from 30 days prior to planting till 11 inch tall corn.
mesotrione @ 0.188 to 0.24 lb/A	Annual broadleaf weeds.	<b>Callisto 4L</b> 6 to 7.7 oz/A.	Preemergence.	<b>Do not plant crops other than corn in treated fields during the same season.</b>
fluthiacet-methyl + mesotrione @ 0.0054 + 0.093 lb/A	Annual broadleaf weeds.	<b>Solstice</b> 3.15 oz/A.	Preemergence.	<b>Do not plant crops other than corn in treated fields during the same season.</b>
metolachlor + mesotrione + atrazine @ 1.3 + 0.168 + 1.3 lb/A	Annual grasses and broadleaf weeds.	<b>Lexar EZ 3.75 SE</b> 3 qt/A.	Preemergence.	<b>Do not plant crops other than corn in treated fields during the same season. Do not exceed 3.5 qt/A per year.</b>
thiencarbazono + isoxaflutole @ 0.019 to 0.03 + 0.05 to 0.08 lb/A	Annual grasses and broadleaf weeds.	<b>Corvus 2.63 SC</b> 3.3 to 5.6 oz/A. See label for soil type restrictions.	Preemergence.	Do not apply to corn treated with Counter, Lorsban or other OP or carbamate insecticides – see label for precautions. Do not plant crops other than corn in treated fields during the same season.
pyroxasulfone @ 0.08 to 0.16 lb/A	Annual grasses, pigweed, glyphosate-resistant ryegrass.	<b>Zidua</b> 1.5 to 3 oz/A.	Preemergence.	Add atrazine for improved control of cocklebur and morningglory. Rates vary based on soil type. Do not apply more than one application to corn in the spring.
pyroxasulfone + fluthiacet-methyl @ 0.11 to 0.16 lb/A + 0.003 to 0.005 lb/A	Annual grasses, pigweed, glyphosate-resistant ryegrass.	<b>Anthem Maxx</b> 2.5 to 5 oz/A.	Preemergence.	Add atrazine for improved control of cocklebur and morningglory. Rates vary based on soil type. Do not apply more than one application to corn in the spring.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
<b>Postemergence</b>				
atrazine @ 2 lb/A	Most small-seeded annuals. More effective on broadleaf weeds, red rice and sicklepod.	<p><b>AAtrex, Atrazine</b> 2.5 lb/A 80W or 2 qt/A of 4L or 2.2 lb/A Nine-0. Select rate according to soil texture. No surfactant recommended on label. Dual, Lasso or Outlook may be added if no soil-applied grass herbicide was used.</p> <p><b>AAtrex, Atrazine + oil</b> 2.5 lb/A 80W or 2 qt/A 4L or 2.2 lb/A Nine-0 + 1 qt/A oil concentrate.</p>	After corn emergence, before grass weeds reach ½ inch or broadleaf 1½ inches.	Do not apply if corn is taller than 12 inches. Do not plant crops other than corn or grain sorghum in treated field until following season. After June 10, do not plant any crop other than corn or grain sorghum the following year. Do not apply more than 2.5 lb/A active atrazine per season.
metolachlor @ 0.95 to 1.9 lb/A	Annual grass and some small-seeded broadleaf weeds.	<b>Dual II Magnum 7.62 EC or Cinch</b> 1 to 2 pt/A.	Apply before weeds emerge or tank mix with a postemergence herbicide like glyphosate or Liberty.	Do not apply more than 3.9 pints of Dual Magnum per acre per year.
2,4-D amine @ 0.5 lb/A	Morningglory, cocklebur and most other young broadleaf weeds.	<b>2,4-D amine</b> 1 pt/A of 4 lb/gal 2,4-D + 0.25% NIS.	Apply when weeds are small and corn is under 12 inches; however, effective results can be obtained with later application.	After corn is more than 12 inches, apply spray directly on weeds with a drop-type nozzle between the corn row and not on the terminal growth of corn. <b>AVOID DRIFT to cotton and soybeans. Follow all State Plant Board Regulations.</b>
dicamba @ 0.25 lb/A	Same as above.	<b>Clarity 4 SL</b> 0.5 pt/A + 0.25% NIS.	From corn emergence up to 15 inches tall.	Ground application only. <b>Drift is extremely toxic to soybeans.</b> Do not apply after soybeans begin to emerge in general area. Less toxic than 2,4-D to cotton. <b>Follow all State Plant Board regulations.</b>
dicamba + diflufenzopyr	Pigweed, morningglory and most other annual broadleaf weeds.	<b>Status</b> 5 to 10 oz/A + 0.25% NIS.	Apply on 4-inch to 36-inch-tall corn.	Status requires an NIS at 0.25% v/v. Do not tank mix with 2,4-D or clopyralid-containing products. Ground application only. <b>Drift is extremely toxic to soybeans. Do not apply after soybeans begin to emerge in general area. Less toxic to cotton than 2,4-D.</b> Follow all State Plant Board regulations.
bentazon @ 0.75 to 1 lb/A	Cocklebur, ragweed, jimsonweed, smartweed, prickly sida, velvetleaf and yellow nutsedge.	<b>Basagran 4 SL</b> 0.75 to 1 qt/A. <b>Can be tank mixed with 0.5 to 0.75 lb/A active atrazine.</b>	Postemergence. See label for specific timing for weed desired. Corn tolerant at all stages.	May be tank mixed with atrazine. See label. Best treatment for smartweed.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
<b>FIELD CORN</b>				
<b>Postemergence [cont.]</b>				
nicosulfuron @ 0.031 lb/A	Johnsongrass, broadleaf signalgrass, foxtail and shattercane.	<b>Accent Q 54.5 DF</b> Accent Q + nonionic surfactant (80%) or crop oil concentrate and 28% or 32% UAN liquid fertilizer (optional). 0.5 oz/A + 2 pt/100 gal or 1 gal/100 gal and 4 gal/100 gal. Tank mix with atrazine for broadleaf weeds.	Apply to 4- to 10-inch seedling and 8- to 12-inch rhizome johnsongrass. If regrowth occurs, apply a second application when johnsongrass is 8 to 10 inches tall. 1- to 2-leaf broadleaf signalgrass. May be applied to 2- to 6-leaf stage of corn.	Repeat application may be required to control regrowth. <b>Do not apply to corn treated with Counter or Counter 20CR insecticide unless IT (Clearfield) corn is planted. See label for restrictions with other organo-phosphate insecticides and postemergence herbicides. Do not apply during cool, cloudy weather.</b>
halosulfuron @ 0.063 lb/A or halosulfuron + thifensulfuron @ 0.031 + 0.004	Nutsedge, cocklebur. See label for tank mixes to broaden weed spectrum.	<b>Halomax, Permit 75 WG, or Permit Plus</b> 1 to 1.33 oz/A for nutsedge. Add a nonionic surfactant or crop oil concentrate. May use two applications not to exceed 2.67 oz/A total rate. Use 0.75 oz Permit Plus.	Postemergence from corn spike through layby. 4- to 12-inch nutsedge 1- to 9-inch cocklebur	See label for mixtures and other precautions. Clean tank with ammonia. Do not use Permit Plus after 5 collars or 6 leaf.
halosulfuron + dicamba @ 0.075 + 0.2 lb/A	Ragweed, horseweed, nutsedge and broadleaf weeds.	<b>Yukon 67.5 DG</b> 6 oz/A.	From corn emergence to 15 inches.	Ground application only. <b>Drift is extremely toxic to soybeans.</b> Do not apply after soybeans begin to emerge in general area. Less toxic than 2,4-D to cotton. <b>Follow all State Plant Board regulations.</b>
rimsulfuron + thifensulfuron @ 0.014 + 0.003 lb/A	Some grass and broadleaf weeds.	<b>Resolve Q 22.4 DG</b> 1.25 oz/A.	Early postemergence to corn V6 stage of growth or earlier, or up to 20-inch corn, whichever is more restrictive.	<b>Some hybrids are sensitive to rimsulfuron. Consult with seed supplier for sensitivity prior to applying.</b>
nicosulfuron + rimsulfuron @ 0.023 + 0.012 lb/A	Annual grass and broadleaf weeds.	<b>Steadfast Q 37.7 DG</b> 1.5 oz/A. Add a crop oil concentrate.	Apply to small weeds. Early post to corn, V6 or up to 20-inch corn, whichever is more restrictive.	<b>Some hybrids are sensitive to rimsulfuron. Consult with seed supplier for sensitivity rating prior to applying.</b> Do not apply to corn treated with Counter or Counter 20 CR insecticides. See label for other precautions.
mesotrione @ 0.094 lb/A	Annual broadleaf weeds.	<b>Callisto 4L</b> 3 oz/A. Add surfactant.	May be applied up to 30 inches or 8-leaf stage of corn for extended morningglory control.	Do not apply to corn treated with Counter or Lorsban insecticides. See label.
thiencarbazone + tembo-trione @ 0.013 + 0.068 lb/A	Annual grass and broadleaf weeds.	<b>Capreno 3.45L</b> 3 oz/A. Add surfactant.	Apply when corn has between 1 and 5 collars.	<b>Some hybrids are sensitive to ALS-inhibiting herbicides. Consult with seed supplier for sensitivity rating prior to use.</b> Do not apply to corn treated with Counter or Lorsban insecticides.
metolachlor + mesotrione + bicyclopyrone + atrazine @ 1.34 + 0.148 + 0.037 + 0.624 lb/A	Grasses and broadleaf weeds.	<b>Acuron</b> 2.5 qt/A.	From corn emergence up to 12 inches.	Do not plant crops other than corn in treated area.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
S-metolachlor + mesotrione + atrazine @ 1.3 + 0.168 + 1.3 lb/A	Annual grasses and broadleaf weeds.	<b>Lexar EZ 3.75 SE</b> 3 qt/A.	Early postemergence.	<b>Do not plant crops other than corn in treated fields during the same season. Do not exceed 3.5 qt/A per year.</b>
thiencarbazone + isoxaflutole @ 0.019 to 0.03 + 0.05 to 0.08 lb/A	Annual grasses and broadleaf weeds.	<b>Corvus 2.63 SC</b> 3.3 to 5.6 oz/A. See label for soil type restrictions.	Early postemergence.	Do not apply to corn treated with Counter, Lorsban or other OP or carbamate insecticides – see label for precautions. Do not plant crops other than corn in treated fields during the same season. The addition of atrazine improves control of pigweed and morningglory.
pyroxasulfone @ 0.08 to 0.21 lb/A	Annual grasses, pigweed, glyphosate-resistant ryegrass.	<b>Zidua</b> 1.5 to 4 oz/A.	Apply before weeds emerge or tank mix with a postemergence herbicide like glyphosate or Liberty. Emergence to V4.	4 oz/A max use rate per application. 5 oz/A total allowed per season.
pyroxasulfone + fluthiacetmethyl @ 0.08 to 0.16 lb/A + 0.003 to 0.005 lb/A	Annual grasses, pigweed, glyphosate-resistant ryegrass.	<b>Anthem Maxx</b> 2.5 to 5 oz/A.	Apply before weeds emerge or tank mix with a postemergence herbicide like glyphosate or Liberty. Emergence to V4.	Do not exceed a maximum of 8.15 oz per season.
mesotrione + rimsulfuron @ 0.078 + 0.019 lb/A	Annual grass and broadleaf weeds.	<b>Realm Q 38.75 DG</b> 4 oz/A. Add surfactant.	Early postemergence to corn V6 stage of growth or up to 20 inches, whichever is more restrictive.	<b>Some hybrids are sensitive to rimsulfuron. Consult with seed supplier for sensitivity rating prior to applying.</b> Do not apply to corn treated with Counter or Counter 20 CR insecticides. See label for other precautions.
topramezone + atrazine @ 0.0164 lb/A + 1 lb/A	Pigweed, horseweed, velvetleaf, morningglory, barnyardgrass, fall panicum and broadleaf signalgrass.	<b>Armezon or Impact + Atrazine</b> 0.75 oz/A + 1 qt/A. COC or MSO at 1% v/v.	Postemergence up to 45 days from harvest.	Apply to corn when weeds are small and actively growing. Use an MSO at 1% v/v and tank mix with atrazine for larger weeds and best results. Can use up to 1 oz Armezon for larger weeds.
topramezone + dimethenamid @ 0.016 lb/A + 0.82 lb/A	Pigweed, horseweed, velvetleaf, morningglory, barnyardgrass, fall panicum and broadleaf signalgrass.	<b>Armezon PRO</b> 20 oz/A.	Emergence through V8 or 30-inch corn.	Apply to corn when weeds are small and actively growing. Use an MSO at 1% v/v and tank mix with atrazine for larger weeds and best results.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
<b>FIELD CORN</b>				
<b>HERBICIDE-TOLERANT CULTIVARS – Check suitability of available hybrids with county agent.</b>				
glyphosate @ 0.75 to 1 lb/A	Most annual grass and broadleaf weeds and Johnsongrass.	<b>Glyphosate (4 lb/gal formulations)</b> 2 pt/A.	May be applied up to 30 inches or 8-leaf stage of corn.	Apply only to Roundup Ready corn. Single in-crop applications not to exceed 1 lb/A and multiple in-crop applications not to exceed 2 lb/A total. See label for tank mixes.
glyphosate + atrazine @ 0.75 to 1 lb/A + 1 lb/A	Same as above plus residual control of broadleaf weeds. Improved morningglory control.	<b>Glyphosate (4 lb/gal formulations) + AAtrex</b> 2 pt/A + 1 qt/A.	Prior to 12-inch corn.	Apply only to Roundup Ready corn. Single in-crop applications not to exceed 1 lb/A and multiple in-crop applications not to exceed 2 lb/A total. See label for tank mixes.
glyphosate + atrazine + thifensulfuron/rimsulfuron @ 0.75 to 1 lb/A + 1 lb/A + 0.014 lb/A	Most annual grass and broadleaf weeds. Improved residual control of annual grass.	<b>Glyphosate (4 lb/gal formulations) + AAtrex + Resolve Q</b> 2 pt/A + 2 pt/A + 1.25 oz/A or <b>Realm Q</b> 2 pt/A + 2 pt/A + 4 oz/A.	Prior to 12-inch corn or 7 collar, whichever comes first.	Apply only to Roundup Ready corn. Single in-crop applications not to exceed 1 lb/A and multiple in-crop applications not to exceed 2 lb/A total. See label for tank mixes. <b>Some hybrids are sensitive to rimsulfuron. Consult with seed supplier for sensitivity rating prior to applying.</b>
glyphosate + mesotrione @ 0.75 to 1 lb/A + 0.094 lb/A	Annual grass and broadleaf weeds with residual activity.	<b>Glyphosate (4 lb/gal formulations) + Callisto 4L</b> 2 pt/A + 3 oz/A.	May be applied up to 30 inches or 8-leaf stage of corn.	Apply only to Roundup Ready corn. Single in-crop applications not to exceed 1 lb/A and multiple in-crop applications not to exceed 2 lb/A total. See label for tank mixes. Do not apply to corn treated with Counter or Lorsban insecticides.
glufosinate @ 0.4 lb/A	Most annual grass and broadleaf weeds.	<b>Glufosinate (280 formulations)</b> 22 oz/A.	May be applied to corn through the V7 growth stage. May be tank mixed with other corn herbicides for residual control.	<b>Apply only to Liberty Link corn varieties or those containing Herculex or Smartstax traits.</b> Do not apply more than 44 ounces of glufosinate per season.
glyphosate + S-metolachlor + mesotrione @ 0.94 + 0.94 + 0.094 lb/A	Most annual grass and broadleaf weeds.	<b>Glyphosate (4 lb/gal formulations) + Dual Magnum II 7.64 EC or Cinch 7.64 EC + Callisto 4L</b> 30 oz + 16 oz + 3 oz/A. or <b>Halex GT</b> 3.6 pt/A.	From corn emergence to 30" or 8-leaf corn.	Tank mix with atrazine will improve residual morningglory control. Sequence at 2.5 to 3.5 pt/A contains glyphosate and metolachlor. The Halex GT rate is 3.6 to 4 pt/A.

Crop, Situation, and Active Chemical Per Broadcast Acre	Weeds Controlled	Formulated Material Per Broadcast Acre	Time of Application	Method of Application and Precautions
<b>Preharvest</b>				
carfentrazone @ 0.0312 lb/A	Morningglory desiccation.	<b>Aim 2 EC</b> 2.0 oz/A. Add 1% crop oil concentrate.	7 days prior to harvest.	Good coverage is critical to Aim activity. 10 gpa is recommended.
sodium chlorate @ 4.5 to 6 lb/A	Desiccation of green vegetation.	<b>Sodium Chlorate</b> Several brands and trade names available. 2 gal of 3 lb/gal or 1 gal of 6 lb/gal.	7 to 10 days prior to harvest.	Use a labeled brand and follow label directions.
<b>Post Harvest</b>				
paraquat + flumioxazin @ 0.625 + 0.063 lb/A	Pigweed.	<b>Paraquat (2 or 3 lb/gal formulations) + Valor 51 WDG</b> 40 or 27 oz/A + 2.0 oz/A. Add 1% COC.	Apply to small pigweed after harvest.	Apply 30 days prior to planting wheat.
paraquat + S-metolachlor @ 0.625 + 0.95 lb/A	Pigweed and annual grass.	<b>Paraquat (2 or 3 lb/gal formulations) + Dual Magnum 7.62 EC</b> 40 or 27 oz/A + 1 pt/A. Add 1% COC.	Apply to small pigweed after harvest.	Apply to acres that <b>will not be</b> planted to small grains (wheat). Follow Dual label on total use rates.
paraquat + metribuzin @ 0.625 + 0.141 lb/A	Volunteer corn, pigweed and other weeds.	<b>Paraquat (2 or 3 lb/gal formulations) + metribuzin 75 DF</b> 40 or 27 oz/A + 3 oz/A. Add 1% COC.	Apply to 6-inch volunteer corn.	If planting wheat, use a metribuzin-tolerant variety.
		<b>OR</b> <b>Boundary</b> 2 pt/A.	3- to 4-inch ryegrass. September through November.	Add gramoxone 48 oz/A. Section 24C.
2,4-D amine @ 0.75 lb/A	Pigweed and other broadleaf weeds.	<b>2,4-D amine</b> 1.5 pt/A of 4 lb/gal 2,4-D.	Apply to 4- to 6-inch pigweed.	Avoid drift to cotton and soybeans. Be aware of state regulations on 2,4-D. 7-day plant-back to wheat. Due to potential off-target movement to maturing soybeans, dicamba is not recommended for use post-harvest in corn.
saflufenacil @ 0.022 to 0.044 lb/A	Pigweed and other broadleaf weeds.	<b>Sharpen 2.85 SC</b> 1.0 oz/A + 1% v/v MSO.	Apply to 4- to 6-inch pigweed.	Avoid off-target drift to soybean.