

CROP REPLANT AND ROTATION GUIDE FOR COTTON HERBICIDES*

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Aim 2 EC	C,CT,S,R,GS All	I 30 days	
Anthem Flex	S,C R SG W CT,P,SF	I 10 months 11 months I 4 months	
Assure II	CT,S All	I 4 months	
Brake FX 32 oz	CT C,GS,R,S All	I 10 months 18 months	Corn rotational interval increases with higher rate.
Caparol/others	CT SG,GS, as cover crops. All	I I† FY	† Must be plowed and not used for food or feed.
Cotoran	CT All	I 6 months	
Direx	CT† All	I FY	† Do not retreat with second application in same year.
Dual Magnum	S,C,CT,GS† SG Rice All	I 4.5 months Next spring 18 months	† Use Concep-treated grain sorghum seed.
Envoke	C,GS,S,R CT W All	7 months 30 days 3 months 18 months	Cotton rotation increases with higher rates.
Fierce	CT C GS R S W	45 days 30 days 12 months 10 months I 1 month	
Fusilade/Fusion	CT,S All	I 2 months	
Glufosinate	CT,C,S,R W GS	I 70 days 180 days	
Glyphosate	All	I	
Linex	C,GS,S All	I 4 months	Thoroughly rework soil before replanting. Do not retreat with second application. Plant corn at least 1.75 inches deep and grain sorghum at least 1 inch deep.
MSMA, DSMA	All	I	Research has shown arsenical herbicides (MSMA/DSMA) can cause straighthead in rice. Precautions for straighthead should be taken when rice is grown following cotton.

Herbicide	Replant/ Crop Rotation	Time Interval	Precautions
Paraquat			No restrictions.
Poast Plus			No restrictions.
Prowl, Pendimax	CT,S W,B All	I 4 months FY	Do not rework soil deeper than treated zone.
Prowl, Pendimax (2X rate)	CT,S All	I FY	Do not rework soil deeper than treated zone.
Reflex//Flexstar/ Cheetah Max/Sinister	S,CT W,SG C,R,GS SF	I 4 months 10 months 18 months	
Select Max, Tapout	C	30 days	
Staple LX	CT S R GS C	I 10 months 9 months Do not rotate the following season. IMI-resistant corn – 9 months. Any other variety – 10 months if not more than 3.8 oz Staple XL was applied.	
Suprend	W CT,C,GS,R,S	3 months 7 months	
Treflan/others	CT,S W,B All	I Fall FY	
Treflan/others (2X rate)	CT,S Rice All	I FY 2 years	
Valor	C,CT,R,GS, W,SF S All others	30 days I 12 months	Must receive 1 inch of rain.
Zidua	CT,C,S W GS R, others	I 70 days 6 months FY	

*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.

Key Crop		Timing
C = Corn	S = Soybeans	All = All crops not specified
CT = Cotton	SF = Sunflowers	I = Immediately
B = Barley	SG = Small Grains	FY = Following year (usually spring)
GS = Grain Sorghum	W = Wheat	
R = Rice		